

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA375401

DDX5 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB,1:500 - 1:2000 IHC,1:50 - 1:200
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-614 of human DDX5 (NP_004387.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	60kDa/69kDa
Gene Name:	DEAD-box helicase 5
Database Link:	<u>Entrez Gene 1655 Human</u> <u>P17844</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


DDX5 Rabbit Polyclonal Antibody – TA375401

Background:DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are
putative RNA helicases. They are implicated in a number of cellular processes involving
alteration of RNA secondary structure, such as translation initiation, nuclear and
mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution
patterns, some members of this family are believed to be involved in embryogenesis,
spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein,
which is a RNA-dependent ATPase, and also a proliferation-associated nuclear antigen,
specifically reacting with the simian virus 40 tumor antigen. Alternative splicing results in
multiple transcript variants.

Synonyms:

DKFZp434E109; DKFZp686J01190; G17P1; HELR; HLR1; HUMP68; p68

Product images:



Western blot analysis of extracts of various cell lines, using DDX5 antibody (TA375401) at 1:2000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 5s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US