

## **Product datasheet for TA375481**

## **DKK1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB.1:500 - 1:1000 Reactivity: Human, Mouse, Rat

**Modifications:** Unmodified

Host: Rabbit Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 71-266 of

human DKK1 (NP\_036374.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 28kDa

Gene Name: dickkopf WNT signaling pathway inhibitor 1

Database Link: Entrez Gene 22943 Human

O94907

Background: This gene encodes a protein that is a member of the dickkopf family. It is a secreted protein

with two cysteine rich regions and is involved in embryonic development through its

inhibition of the WNT signaling pathway. Elevated levels of DKK1 in bone marrow plasma and peripheral blood is associated with the presence of osteolytic bone lesions in patients with

multiple myeloma.

Synonyms: Dickkopf-1; DKK-1; hDkk-1; SK




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of various cell lines, using DKK1 antibody (TA375481) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.