

Product datasheet for TA375490S

PSD93 (DLG2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Phospho Y340

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A phospho synthetic peptide corresponding to residues surrounding Y340 of human

PSD93/chapsyn-110/DLG2.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Gene Name: discs large MAGUK scaffold protein 2

Database Link: Entrez Gene 1740 Human

Q15700

Background: This gene encodes a member of the membrane-associated guanylate kinase (MAGUK) family.

The encoded protein forms a heterodimer with a related family member that may interact at postsynaptic sites to form a multimeric scaffold for the clustering of receptors, ion channels, and associated signaling proteins. Multiple transcript variants encoding different isoforms have been found for this gene. Additional transcript variants have been described, but their

full-length nature is not known. [provided by RefSeq, Dec 2008]

Synonyms: chapsyn-110; DKFZp781D1854; DKFZp781E0954; FLJ37266; MGC131811;

OTTHUMP00000165971; PSD-93; PSD93




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using Phospho-PSD93/chapsyn-110/DLG2-Y340 antibody ([TA375490]) at 1:1000 dilution.293T cells were treated by ATP(5 mM) at 30°C for 1 hour.NIH/3T3 cells were treated by ATP(5 mM) at 30°C for 1 hour.C6 cells were treated by ATP(5 mM) at 30°C for 1 hour.|Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% BSA.|Detection: ECL Basic Kit.|Exposure time: 1s.