

## **Product datasheet for TA375601**

## Product data:

**Product Type:** Primary Antibodies

**DPP9 Rabbit Polyclonal Antibody** 

Applications: WB

Recommended Dilution: WB,1:200 - 1:2000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-100 of

human DPP9 (NP\_631898.3).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size:95kDa/98kDa/101kDaGene Name:dipeptidyl peptidase 9

Database Link: Entrez Gene 91039 Human

Q86TI2

**Background:** This gene encodes a protein that is a member of the S9B family in clan SC of the serine

proteases. The protein has been shown to have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. Although the activity of this protein is similar to that of dipeptidyl peptidase 4 (DPP4), it does not appear to be membrane bound. In general, dipeptidyl peptidases appear to be involved in the regulation of the activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. Several transcript variants of this gene have been described

but not fully characterized.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

DKFZp762F117; DP9; DPLP9; DPRP-2; DPRP2; FLJ16073

## **Product images:**



Western blot analysis of extracts of various cell lines, using DPP9 antibody (TA375601) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10S.