

Product datasheet for TA375639

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Dystrobrevin alpha (DTNA) Rabbit Polyclonal Antibody

Human, Mouse

Product data:

Reactivity:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 240-480 of

human DTNA (NP_001381.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 22kDa/36kDa/41-44kDa/50kDa/58-83kDa

Gene Name: dystrobrevin alpha

Database Link: Entrez Gene 1837 Human

Q9Y4J8

Background: The protein encoded by this gene belongs to the dystrobrevin subfamily of the dystrophin

family. This protein is a component of the dystrophin-associated protein complex (DPC), which consists of dystrophin and several integral and peripheral membrane proteins, including dystroglycans, sarcoglycans, syntrophins and alpha- and beta-dystrobrevin. The DPC localizes to the sarcolemma and its disruption is associated with various forms of

muscular dystrophy. Mutations in this gene are associated with left ventricular

noncompaction with congenital heart defects. Multiple alternatively spliced transcript

variants encoding different isoforms have been identified for this gene.






Synonyms:

D18S892E; DRP3; DTN; DTN-1; DTN-2; DTN-3; Dystrobrevin-alpha; FLJ96209; LVNC1; OTTHUMP00000163152; OTTHUMP00000163155

Product images:



Western blot analysis of extracts of various cell lines, using DTNA antibody (TA375639) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.