

Product datasheet for TA375674

DYNLT1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB,1:200 - 1:2000 Reactivity:

Modifications: Unmodified

Host: Rabbit

Isotype: **IgG**

Clonality: Polyclonal

Recombinant fusion protein containing a sequence corresponding to amino acids 1-113 of Immunogen:

human DYNLT1 (NP_006510.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Human, Mouse

Shelf life: one year from despatch. Stability:

Predicted Protein Size: 12kDa

Gene Name: dynein light chain Tctex-type 1

Database Link: Entrez Gene 6993 Human

P63172



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes a component of the motor complex, cytoplasmic dynein, which transports cellular cargo along microtubules in the cell. The encoded protein regulates the length of primary cilia which are sensory organelles found on the surface of cells. The protein encoded by this gene interacts with viral proteins, like the minor capsid protein L2 of human papillomavirus, and is required for dynein-mediated delivery of the viral nucleic acid to the host nucleus. This protein interacts with oncogenic nucleoporins to disrupt gene regulation and cause leukemic transformation. Pseudogenes of this gene are present on chromosomes 4 and 17. Alternative splicing results in multiple transcript variants encoding different isoforms.

Synonyms: CW-1; MGC111571; TCTEL1; TCTEX-1; TCTEX1

Product images:



Western blot analysis of extracts of various cell lines, using DYNLT1 antibody (TA375674) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 60s.