

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA375697

ECH1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB,1:500 - 1:2000 IHC,1:50 - 1:200
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-328 of human ECH1 (NP_001389.2).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	35kDa
Gene Name:	enoyl-CoA hydratase 1, peroxisomal
Database Link:	<u>Entrez Gene 1891 Human</u> <u>Q13011</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


CRIGENE ECH1 Rabbit Polyclonal Antibody – TA375697

HPXEL

Background:This gene encodes a member of the hydratase/isomerase superfamily. The gene product
shows high sequence similarity to enoyl-coenzyme A (CoA) hydratases of several species,
particularly within a conserved domain characteristic of these proteins. The encoded protein,
which contains a C-terminal peroxisomal targeting sequence, localizes to the peroxisome.
The rat ortholog, which localizes to the matrix of both the peroxisome and mitochondria, can
isomerize 3-trans,5-cis-dienoyl-CoA to 2-trans,4-trans-dienoyl-CoA, indicating that it is a
delta3,5-delta2,4-dienoyl-CoA isomerase. This enzyme functions in the auxiliary step of the
fatty acid beta-oxidation pathway. Expression of the rat gene is induced by peroxisome
proliferators.

Synonyms:

Product images:



Western blot analysis of extracts of various cell lines, using ECH1 Antibody (TA375697) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 10s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US