

## **Product datasheet for TA375737**

## Product datasileet for 1A3/3/3/

## **Ephrin B2 (EFNB2) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 28-229 of

human Ephrin B2 (NP\_004084.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 36kDa

Gene Name: ephrin B2

Database Link: P52799

**Background:** This gene encodes a member of the ephrin (EPH) family. The ephrins and EPH-related

receptors comprise the largest subfamily of receptor protein-tyrosine kinases and have been

implicated in mediating developmental events, especially in the nervous system and in

erythropoiesis. Based on their structures and sequence relationships, ephrins are divided into

the ephrin-A (EFNA) class, which are anchored to the membrane by a

glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are

transmembrane proteins. This gene encodes an EFNB class ephrin which binds to the EPHB4

and EPHA3 receptors.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ephrin-B2; EPLG5; HTK-L; HTKL; LERK-5; LERK5; MGC126226; MGC126227; MGC126228

## **Product images:**



Western blot analysis of extracts of various cell lines, using Ephrin B2 antibody (TA375737) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.