

Product datasheet for TA375794S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EIF2B1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF/ICC,1:50 - 1:200

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 34kDa

Gene Name: eukaryotic translation initiation factor 2B subunit alpha

Database Link: Entrez Gene 1967 Human

Q14232

Background: This gene encodes one of five subunits of eukaryotic translation initiation factor 2B (EIF2B), a

GTP exchange factor for eukaryotic initiation factor 2 and an essential regulator for protein synthesis. Mutations in this gene and the genes encoding other EIF2B subunits have been


associated with leukoencephalopathy with vanishing white matter.


Synonyms: EIF-2B; EIF-2Balpha; EIF2B; EIF2BA; MGC117409; MGC125868; MGC125869





Product images:





Western blot analysis of various lysates using EIF2B1 Rabbit pAb ([TA375794]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.

Immunofluorescence analysis of U2OS cells using EIF2B1 Rabbit pAb ([TA375794]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.