

Product datasheet for TA375811

EIF3S4 (EIF3G) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:200 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-320 of

human EIF3G (NP_003746.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 35kDa

Gene Name: eukaryotic translation initiation factor 3 subunit G

Database Link: Entrez Gene 8666 Human

O75821

Background: This gene encodes a core subunit of the eukaryotic translation initiation factor 3 (eIF3)

complex, which is required for initiation of protein translation. An N-terminal caspase cleavage product of the encoded protein may stimulate degradation of DNA. A mutation in

this gene is associated with narcolepsy. [provided by RefSeq, Jul 2016]

Synonyms: eIF-3-delta; eIF3-delta; EIF3-P42; eIF3-p44; EIF3S4




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using EIF3G antibody (TA375811) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 1s.



Immunofluorescence analysis of U-2 OS cells using EIF3G Polyclonal Antibody (TA375811) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.