

## **Product datasheet for TA375905**

## **ENPP3 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 646-875 of

human ENPP3 (NP\_005012.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 100kDa

**Gene Name:** ectonucleotide pyrophosphatase/phosphodiesterase 3

Database Link: Entrez Gene 5169 Human

<u>014638</u>

**Background:** The protein encoded by this gene belongs to a series of ectoenzymes that are involved in

hydrolysis of extracellular nucleotides. These ectoenzymes possess ATPase and ATP

pyrophosphatase activities and are type II transmembrane proteins. Expression of the related rat mRNA has been found in a subset of immature glial cells and in the alimentary tract. The corresponding rat protein has been detected in the pancreas, small intestine, colon, and liver.

The human mRNA is expressed in glioma cells, prostate, and uterus. Expression of the human protein has been detected in uterus, basophils, and mast cells. Two transcript

variants, one protein coding and the other non-protein coding, have been found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

B10; CD203c; gp130RB13-6; NPP3; OTTHUMP00000017901; PD-IBETA; PDNP3

## **Product images:**



Western blot analysis of extracts of various cell lines, using ENPP3 Rabbit pAb (TA375905) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 1s.