

Product datasheet for TA375930

Eph receptor A3 (EPHA3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: WB Recommended Dilution: WB.1:500 - 1:2000 **Reactivity:** Human **Modifications:** Phospho Y779 Host: Rabbit Isotype: lgG **Clonality:** Polyclonal A synthetic phosphorylated peptide around Y779 of human EphA3 (NP_005224.2). Immunogen: Formulation: Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. **Concentration:** lot specific **Purification:** Affinity purification Unconjugated **Conjugation:** Storage: Store at -20°C. Avoid freeze / thaw cycles. Shelf life: one year from despatch. Stability: Predicted Protein Size: 60kDa/110kDa Gene Name: **EPH receptor A3** Database Link: Entrez Gene 2042 Human P29320 **Background:** This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene encodes a protein that binds ephrin-A ligands. Two alternatively spliced transcript



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

variants have been described for this gene.


OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Synonyms:

ETK; ETK1; HEK; HEK4; TYRO4

Product images:



Western blot analysis of extracts of 293T and Hela cells, using Phospho-EphA3-Y779 antibody (TA375930) at 1:2000 dilution. 293T cells were treated by Pervanadate (1mM) for 30 minutes after serum-starvation overnight.HeLa cells were treated by Pervanadate (1nM) for 30 minutes after serum-starvation overnight. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% BSA.|Detection: ECL Basic Kit.|Exposure time: 1s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US