

## **Product datasheet for TA376033S**

## **EXOC5 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:1000 - 1:2000

IHC,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 340-610 of

human EXOC5 (NP\_006535.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 81kDa

**Gene Name:** exocyst complex component 5

Database Link: Entrez Gene 10640 Human

<u>000471</u>

**Background:** The protein encoded by this gene is a component of the exocyst complex, a multiple protein

complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also

essential for the biogenesis of epithelial cell surface polarity.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

DKFZp666H126; hSec10; PRO1912; SEC10; SEC10L1; SEC10P

## **Product images:**



Western blot analysis of extracts of various cell lines, using EXOC5 antibody ([TA376033]) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.