

Product datasheet for TA376074S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

BLBP (FABP7) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, WB

Recommended Dilution: WB,1:100 - 1:500

IF/ICC,1:50 - 1:200

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 15kDa

Gene Name: fatty acid binding protein 7

Database Link: Entrez Gene 2173 Human

O15540

Background: The gene encodes a small, highly conserved cytoplasmic protein that bind long-chain fatty

acids and other hydrophobic ligands. The encoded protein is important in the establishment of the radial glial fiber in the developing brain. Alternative splicing and promoter usage results in multiple transcript variants encoding different isoforms. Pseudogenes of this gene


are found on multiple chromosomes.

Synonyms: B-FABP; BLBP; DKFZp547J2313; FABPB; MRG






Product images:




Western blot analysis of various lysates using FABP7 Rabbit pAb ([TA376074]) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 180s.



Immunofluorescence analysis of C6 cells using FABP7 Rabbit pAb ([TA376074]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of U-251 MG cells using FABP7 Rabbit pAb ([TA376074]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.