

Product datasheet for **TA376076**

Fadd Rabbit Polyclonal Antibody


Product data:

| | |
|-----------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | ICC/IF, IP, WB |
| Recommended Dilution: | WB,1:200 - 1:2000 IF,1:50 - 1:200 IP,1:50 - 1:200 |
| Reactivity: | Human, Mouse, Rat |
| Modifications: | Unmodified |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Recombinant protein of human FADD. |
| Formulation: | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Shelf life: one year from despatch. |
| Gene Name: | Fas (TNFRSF6)-associated via death domain |
| Database Link: | Entrez Gene 14082 Mouse Q61160 |
| Background: | Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95 or TNFR-1 receptors). The resulting aggregate called the death-inducing signaling complex (DISC performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling. |
| Synonyms: | GIG3; MGC8528; MORT1 |




[View online »](#)


Product images:



Western blot analysis of extracts of Mouse heart, using FADD antibody (TA376076) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 10s.



Western blot analysis of extracts of Jurkat cells, using FADD antibody (TA376076) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 180s.



Immunoprecipitation analysis of 600ug extracts of Mouse heart cells using 3ug FADD antibody (TA376076). Western blot was performed from the immunoprecipitate using FADD antibody (TA376076) at a dilution of 1:1000.