

# **Product datasheet for TA376125**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Fibroblast activation protein, alpha (FAP) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications:

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 26-280 of

human fibroblast activation protein-α (Fibroblast activation protein-α (FAP)) (NP\_004451.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 26kDa/87kDa

**Gene Name:** fibroblast activation protein alpha

**Database Link:** Entrez Gene 2191 Human

Q12884






#### Background:

The protein encoded by this gene is a homodimeric integral membrane gelatinase belonging to the serine protease family. It is selectively expressed in reactive stromal fibroblasts of epithelial cancers, granulation tissue of healing wounds, and malignant cells of bone and soft tissue sarcomas. This protein is thought to be involved in the control of fibroblast growth or epithelial-mesenchymal interactions during development, tissue repair, and epithelial carcinogenesis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

**Synonyms:** DKFZp686G13158; DPPIV; FAPA; seprase

## **Product images:**



Western blot analysis of extracts of Rat skeletal muscle, using Fibroblast activation protein- $\alpha$  (FAP) antibody (TA376125) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 180s.