

## **Product datasheet for TA376194**

**FCRL3 Rabbit Polyclonal Antibody** 

## Product data:

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:100 - 1:200

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 18-180 of

human FCRL3 (NP\_443171.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 21kDa/22kDa/69kDa/77kDa/80kDa/81kDa

**Gene Name:** Fc receptor like 3

Database Link: Entrez Gene 115352 Human

Q96P31

**Background:** This gene encodes a member of the immunoglobulin receptor superfamily and is one of

several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein contains immunoreceptor-tyrosine activation motifs and immunoreceptor-tyrosine inhibitory motifs in its cytoplasmic domain and may play a role in regulation of the immune system. Mutations in this gene have been associated with rheumatoid arthritis, autoimmune thyroid disease, and systemic lupus erythematosus. Alternative splicing results

in multiple transcript variants.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

FCRH3; hIFGP3; IFGP3; IRTA3; SPAP2; SPAP2a; SPAP2b; SPAP2c; SPAP2d; SPAP2e

## **Product images:**



Western blot analysis of extracts of various cell lines, using FCRL3 antibody (TA376194) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.