

Product datasheet for TA376253S

FIG4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:200 - 1:2000

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 608-907 of

human FIG4 (NP_055660.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 103kDa

Gene Name: FIG4 phosphoinositide 5-phosphatase

Database Link: Entrez Gene 9896 Human

Q92562

Background: The protein encoded by this gene belongs to the SAC domain-containing protein gene family.

The SAC domain, approximately 400 amino acids in length and consisting of seven conserved

motifs, has been shown to possess phosphoinositide phosphatase activity. The yeast homolog, Sac1p, is involved in the regulation of various phosphoinositides, and affects diverse cellular functions such as actin cytoskeleton organization, Golgi function, and maintenance of vacuole morphology. Membrane-bound phosphoinositides function as signaling molecules and play a key role in vesicle trafficking in eukaryotic cells. Mutations in

this gene have been associated with Charcot-Marie-Tooth disease, type 4J.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ALS11; dJ249I4.1; hSac3; KIAA0274; SAC3

Product images:



Western blot analysis of extracts of various cell lines, using FIG4 Rabbit pAb ([TA376253]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.