

Product datasheet for TA376270S

FLCN Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-130 of

human FLCN (NP_653207.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 20kDa/37kDa/64kDa

Gene Name: folliculin

Database Link: Entrez Gene 201163 Human

Q8NFG4

Background: This gene is located within the Smith-Magenis syndrome region on chromosome 17.

Mutations in this gene are associated with Birt-Hogg-Dube syndrome, which is characterized by fibrofolliculomas, renal tumors, lung cysts, and pneumothorax. Alternative splicing of this

gene results in two transcript variants encoding different isoforms.

Synonyms: BHD; DKFZp547A118; FLCL; FLJ45004; FLJ99377; folliculin; MGC17998; MGC23445




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:







Western blot analysis of extracts of various cell lines, using FLCN antibody ([TA376270]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 1s.

Immunofluorescence analysis of C6 cells using FLCN antibody ([TA376270]) at dilution of 1:100. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of NIH/3T3 cells using FLCN antibody ([TA376270]) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using FLCN antibody ([TA376270]) at dilution of 1:100. Blue: DAPI for nuclear staining.