

# **Product datasheet for TA376298**

### **FMO5 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-285 of

human FMO5 (NP\_001138302.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 32kDa/52kDa/60kDa

**Gene Name:** flavin containing monooxygenase 5

Database Link: Entrez Gene 2330 Human

P49326



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:


Metabolic N-oxidation of the diet-derived amino-trimethylamine (TMA) is mediated by flavin-containing monooxygenase and is subject to an inherited FMO3 polymorphism in man resulting in a small subpopulation with reduced TMA N-oxidation capacity resulting in fish odor syndrome Trimethylaminuria. Three forms of the enzyme, FMO1 found in fetal liver, FMO2 found in adult liver, and FMO3 are encoded by genes clustered in the 1q23-q25 region. Flavin-containing monooxygenases are NADPH-dependent flavoenzymes that catalyzes the oxidation of soft nucleophilic heteroatom centers in drugs, pesticides, and xenobiotics. Alternative splicing results in multiple transcript variants.

Synonyms: FMO5

# **Product images:**



Western blot analysis of extracts of various cell lines, using FMO5 antibody (TA376298) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.



Immunofluorescence analysis of U2OS cells using FMO5 antibody (TA376298) at dilution of 1:100. Blue: DAPI for nuclear staining.