

## **Product datasheet for TA376320**

## Fos B (FOSB) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 1-100 of human FOSB

(NP\_006723.2).

**Formulation:** PBS with 0.05% proclin300,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 15kDa/20kDa/26kDa/27kDa/30kDa/31kDa/35kDa

**Gene Name:** FosB proto-oncogene, AP-1 transcription factor subunit

Database Link: Entrez Gene 2354 Human

P53539

**Background:** The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes

encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby

forming the transcription factor complex AP-1. As such, the FOS proteins have been

implicated as regulators of cell proliferation, differentiation, and transformation. Alternatively

spliced transcript variants encoding different isoforms have been found for this gene.

**Synonyms:** AP-1; DKFZp686C0818; G0S3; GOS3; GOSB; MGC42291




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of 293T cells, using FOSB antibody (TA376320) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 1s.



Western blot analysis of extracts of various cell lines, using FOSB antibody (TA376320) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: