

## **Product datasheet for TA376436**

## GABPA Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human, Rat
Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 1-100 of human GABPA

(NP 002031.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 51kDa

**Gene Name:** GA binding protein transcription factor alpha subunit

Database Link: Entrez Gene 2551 Human

Q06546

**Background:** This gene encodes one of three GA-binding protein transcription factor subunits which

functions as a DNA-binding subunit. Since this subunit shares identity with a subunit

encoding the nuclear respiratory factor 2 gene, it is likely involved in activation of cytochrome oxidase expression and nuclear control of mitochondrial function. This subunit also shares identity with a subunit constituting the transcription factor E4TF1, responsible for expression of the adenovirus E4 gene. Because of its chromosomal localization and ability to form heterodimers with other polypeptides, this gene may play a role in the Down Syndrome

phenotype. Two transcript variants encoding the same protein have been found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

E4TF1-60; E4TF1A; NFT2; NRF2; NRF2A; OTTHUMP00000096114

## **Product images:**



Western blot analysis of extracts of various cell lines, using GABPA antibody (TA376436) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 1s.