

## **Product datasheet for TA376468**

**GALNT3** Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of

human GALNT3 (NP 004473.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 22kDa/72kDa

**Gene Name:** polypeptide N-acetylgalactosaminyltransferase 3

**Database Link:** Entrez Gene 2591 Human

Q14435

**Background:** This gene encodes UDP-GalNAc transferase 3, a member of the GalNAc-transferases family.

This family transfers an N-acetyl galactosamine to the hydroxyl group of a serine or threonine residue in the first step of O-linked oligosaccharide biosynthesis. Individual GalNAc-transferases have distinct activities and initiation of O-glycosylation is regulated by a repertoire of GalNAc-transferases. The protein encoded by this gene is highly homologous to

other family members, however the enzymes have different substrate specificities.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

DKFZp686C10199; GalNAc-T3; HFTC; HHS; MGC61909

## **Product images:**



Western blot analysis of extracts of various cell lines, using GALNT3 antibody (TA376468) at 1:1000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 1s.



Immunofluorescence analysis of U-2 OS cells using GALNT3 Rabbit pAb (TA376468) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.