

Product datasheet for TA376519

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glucokinase (GCK) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Reactivity: WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of

human Glucokinase (Glucokinase (GCK)) (NP_000153.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 52kDa

Gene Name: glucokinase

Database Link: Entrez Gene 2645 Human

P35557

Background: Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most

glucose metabolism pathways. Alternative splicing of this gene results in three tissue-specific forms of glucokinase, one found in pancreatic islet beta cells and two found in liver. The protein localizes to the outer membrane of mitochondria. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose-6-phosphate but remains active while glucose is abundant. Mutations in this gene have been associated with non-insulin dependent diabetes mellitus (NIDDM), maturity onset diabetes of the young, type 2

(MODY2) and persistent hyperinsulinemic hypoglycemia of infancy (PHHI).






Synonyms:

GK; GLK; glucokinase; Hexokinase-4; Hexokinase-D; HHF3; HK4; HKIV; HXKP; MODY2

Product images:



Western blot analysis of extracts of various cell lines, using Glucokinase (Glucokinase (GCK)) antibody (TA376519) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 60s.