

## **Product datasheet for TA376565**

## **GIGYF2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** ELISA, WB

**Recommended Dilution:** WB,1:500 - 1:1000

ELISA,Recommended starting concentration is 1 µg/mL. Please optimize the concentration

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn

based on your specific assay requirements.

**Reactivity:** Human, Mouse, Rat

**Modifications: Unmodified** 

Rabbit Host:

IqG Isotype:

Polyclonal Clonality:

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. Formulation:

**Concentration:** lot specific

Affinity purification **Purification:** 

Unconjugated Conjugation:

Store at -20°C. Avoid freeze / thaw cycles. Storage:

Shelf life: one year from despatch. Stability:

**Predicted Protein Size:** 150kDa

GRB10 interacting GYF protein 2 Gene Name:

Entrez Gene 26058 Human Database Link:

Q6Y7W6






**Background:** 

This gene contains CAG trinucleotide repeats and encodes a protein containing several stretches of polyglutamine residues. The encoded protein may be involved in the regulation of tyrosine kinase receptor signaling. This gene is located in a chromosomal region that was genetically linked to Parkinson disease type 11, and mutations in this gene were thought to be causative for this disease. However, more recent studies in different populations have been unable to replicate this association. Alternative splicing results in multiple transcript variants.

Synonyms:

DKFZp686I15154; DKFZp686J17223; FLJ23368; GYF2; KIAA0642; OTTHUMP00000203484; PARK11; PERQ2; PERQ3; TNRC15

## **Product images:**



Western blot analysis of various lysates using GIGYF2 Rabbit pAb (TA376565) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit . Exposure time: 10s.