

Product datasheet for TA376783S

NMDAR1 (GRIN1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB,1:500 - 1:2000

Reactivity: Human, Mouse, Rat

Modifications: Phospho S896/S897

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic phosphorylated peptide around of human Phospho-NMDAR1-S896/S897.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 99kDa/101kDa/103kDa/105kDa/106kDa/107kDa

Gene Name: glutamate ionotropic receptor NMDA type subunit 1

Database Link: Entrez Gene 2902 Human

Q05586

Background: The protein encoded by this gene is a critical subunit of N-methyl-D-aspartate receptors,

members of the glutamate receptor channel superfamily which are heteromeric protein complexes with multiple subunits arranged to form a ligand-gated ion channel. These subunits play a key role in the plasticity of synapses, which is believed to underlie memory and learning. Cell-specific factors are thought to control expression of different isoforms, possibly contributing to the functional diversity of the subunits. Alternatively spliced

transcript variants have been described.

Synonyms: NMDA1; NMDAR1; NR1




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of HeLa cells, using Phospho-NMDAR1-S896/S897 antibody ([TA376783]) at 1:1000 dilution. HeLa cells were treated by UV at room temperature for 30 minutes and PMA (200 nM) at 37°C for 30 minutes after serum-starvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA. | Detection: ECL Basic Kit. | Exposure time: 90s.