

## **Product datasheet for TA376902**

## H3FA (HIST1H3A) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Human, Mouse, Rat, Other (Wide Range)

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide of human Lactic acid-Histone H3-K27.

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 15kDa

Gene Name: histone cluster 1, H3a

Database Link: Entrez Gene 8350 Human

P68431

**Background:** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large

histone gene cluster on chromosome 6p22-p21.3.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

H3/A; H3/b; H3/c; H3/d; H3/f; H3/h; H3/j; H3/k; H3/l; H3FA; H3FB; H3FC; H3FD; H3FF; H3FH; H3FI; H3FJ; H3FK; H3FL

## **Product images:**



Western blot analysis of extracts of C6 cells, using Lactic acid-Histone H3-K27 antibody (TA376902) at 1:1000 dilution.C6 cells were treated by Sodium lactate(100mM) for 24h. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Enhanced Kit. |Exposure time: 180s.