

## Product datasheet for **TA376913**


### HAGH Rabbit Polyclonal Antibody

#### Product data:


Product Type:	Primary Antibodies
Applications:	ICC/IF, WB
Recommended Dilution:	WB,1:500 - 1:2000 IF,1:10 - 1:100
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human HAGH (NP_001035517.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	25kDa/28kDa/33kDa
Gene Name:	hydroxyacylglutathione hydrolase
Database Link:	<a href="#">Entrez Gene 3029 Human Q16775</a>
Background:	The enzyme encoded by this gene is classified as a thiolesterase and is responsible for the hydrolysis of S-lactoyl-glutathione to reduced glutathione and D-lactate. Three transcript variants encoding different isoforms have been found for this gene.
Synonyms:	GLO2; GLX2; GLXII; HAGH1



[View online »](#)

**Product images:**


Western blot analysis of extracts of various cell lines, using HAGH antibody (TA376913) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 1s.



Immunofluorescence analysis of A549 cells using HAGH antibody (TA376913). Blue: DAPI for nuclear staining.