

Product datasheet for TA376923

Product datasileet for TAS/0923

HARS2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Reactivity: WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 217-506 of

human HARS2 (NP_036340.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 54kDa/56kDa

Gene Name: histidyl-tRNA synthetase 2, mitochondrial

Database Link: Entrez Gene 23438 Human

P49590

Background: Aminoacyl-tRNA synthetases are a class of enzymes that charge tRNAs with their cognate

amino acids. The protein encoded by this gene is an enzyme belonging to the class II family of

aminoacyl-tRNA synthetases. Functioning in the synthesis of histidyl-transfer RNA, the

enzyme plays an accessory role in the regulation of protein biosynthesis. The gene is located in a head-to-head orientation with HARS on chromosome five, where the homologous genes

likely share a bidirectional promoter. Mutations in this gene are associated with the pathogenesis of Perrault syndrome, which involves ovarian dysgenesis and sensorineural

hearing loss. Alternative splicing results in multiple transcript variants of this gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

HARS-related; HARSL; HARSR; HisRS; HO3

Product images:



Western blot analysis of extracts of various cell lines, using HARS2 antibody (TA376923) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 5s.