

Product datasheet for TA377006

Product data:

Product Type: Primary Antibodies

HEY2 Rabbit Polyclonal Antibody

Applications: WE

Reactivity: WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 1-100 of human HEY2

(NP 036391.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 35kDa

Gene Name: hes related family bHLH transcription factor with YRPW motif 2

Database Link: Entrez Gene 23493 Human

Q9UBP5

Background: This gene encodes a member of the hairy and enhancer of split-related (HESR) family of basic

helix-loop-helix (bHLH)-type transcription factors. The encoded protein forms homo- or hetero-dimers that localize to the nucleus and interact with a histone deacetylase complex to repress transcription. Expression of this gene is induced by the Notch signal transduction

pathway. Two similar and redundant genes in mouse are required for embryonic

cardiovascular development, and are also implicated in neurogenesis and somitogenesis. Alternatively spliced transcript variants have been found, but their biological validity has not

been determined.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

bHLHb32; CHF1; GRIDLOCK; GRL; hCHF1; HERP; HERP1; HESR-2; HESR2; hHRT2; HRT2; MGC10720

Product images:



Western blot analysis of extracts of various cell lines, using HEY2 antibody (TA377006) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 60s.