

## **Product datasheet for TA377033**

## Product data:

**Product Type:** Primary Antibodies

**HIRA Rabbit Polyclonal Antibody** 

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

**Clonality:** Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 250-550 of

human HIRA (NP\_003316.3).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 89kDa/111kDa

Gene Name: histone cell cycle regulator

Database Link: Entrez Gene 7290 Human

P54198

**Background:** This gene encodes a histone chaperone that preferentially places the variant histone H3.3 in

nucleosomes. Orthologs of this gene in yeast, flies, and plants are necessary for the

formation of transcriptionally silent heterochomatin. This gene plays an important role in the formation of the senescence-associated heterochromatin foci. These foci likely mediate the irreversible cell cycle changes that occur in senescent cells. It is considered the primary candidate gene in some haploinsufficiency syndromes such as DiGeorge syndrome, and

insufficient production of the gene may disrupt normal embryonic development.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

DGCR1; HIR; TUP1; TUPLE1

## **Product images:**



Western blot analysis of extracts of various cell lines, using HIRA Antibody (TA377033) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.