

Product datasheet for TA377047S

Histone H2A.X (H2AFX) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:1000

IHC,1:50 - 1:100 IF,1:50 - 1:100

Reactivity: Human, Mouse, Rat, Other (Wide Range)

Modifications: Unmodified

Host: Rabbit

Isotype: lgG

Clonality: Polyclonal

Immunogen: A synthetic peptide of human Histone H2AX.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 15kDa

Gene Name: H2A histone family member X

Database Link: Entrez Gene 3014 Human

P16104



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stemloop termination motif, and the polyA addition motif.

Synonyms: H2A.X; H2A/X; H2AX

Product images:



Western blot analysis of extracts of various cell lines, using Histone H2AX antibody ([TA377047]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.



Confocal immunofluorescence analysis of HeLa cells using Histone H2AX Polyclonal Antibody ([TA377047]) at dilution of 1:200. Blue: DAPI for nuclear staining.