

## Product datasheet for **TA377055S**

### **H3FT (HIST3H3) Rabbit Polyclonal Antibody**

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	ICC/IF, WB
Recommended Dilution:	WB,1:500 - 1:2000 IF,1:50 - 1:100
Reactivity:	Human, Mouse, Rat, Other (Wide Range)
Modifications:	Acetyl K36
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthetic acetylated peptide around K36 of human Histone H3 (NP_003484.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	15kDa
Gene Name:	histone cluster 3, H3
Database Link:	<a href="#">Entrez Gene 8290 Human Q16695</a>





[View online »](#)


**Background:**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

**Synonyms:**

H3.4; H3/g; H3/t; H3FT; H3t; MGC126886; MGC126888

**Product images:**




Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K36 antibody ([TA377055]) at dilution of 1:100. NIH/3T3 cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.