

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA377104

HLA-DPB1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB,1:500 - 1:2000 IHC,1:50 - 1:200
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 30-225 of human HLA-DPB1 (NP_002112.3).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	29kDa
Gene Name:	major histocompatibility complex, class II, DP beta 1
Database Link:	<u>Entrez Gene 3115 Human</u> <u>P04440</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE HLA-DPB1 Rabbit Polyclonal Antibody – TA377104

Background:HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a
heterodimer consisting of an alpha (DPA) and a beta chain (DPB), both anchored in the
membrane. It plays a central role in the immune system by presenting peptides derived from
extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B
lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and
its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the
two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes
the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain
the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different
molecules.

Product images:



Western blot analysis of extracts of various cell lines, using HLA-DPB1 antibody (TA377104) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 1s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US