

## **Product datasheet for TA377185**

## **HOXA7 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB.1:1000 - 1:2000 Reactivity: Human, Mouse, Rat

**Modifications:** Unmodified

Host: Rabbit Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of

human HOXA7 (NP\_008827.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 25kDa

Gene Name: homeobox A7

**Database Link:** Entrez Gene 3204 Human

P31268

Background: In vertebrates, the genes encoding the class of transcription factors called homeobox genes

> are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome 7 and encodes a DNA-binding transcription factor which may regulate gene expression, morphogenesis, and differentiation. For example, the encoded protein represses the transcription of differentiation-specific genes during

> keratinocyte proliferation, but this repression is then overcome by differentiation signals. This

gene is highly similar to the antennapedia (Antp) gene of Drosophila.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ANTP; Hox-1A; HOX1; HOX1.1; HOX1A; OTTHUMP00000024835

## **Product images:**



Western blot analysis of extracts of various cell lines, using HOXA7 antibody (TA377185) at 1:3000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 90s.