

## **Product datasheet for TA377226**

## **HS3ST3A1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 44-140 of

human HS3ST3A1 (NP\_006033.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 44kDa

**Gene Name:** heparan sulfate-glucosamine 3-sulfotransferase 3A1

Database Link: Entrez Gene 9955 Human

Q9Y663

**Background:** Heparan sulfate biosynthetic enzymes are key components in generating a myriad of distinct

heparan sulfate fine structures that carry out multiple biologic activities. The enzyme encoded by this gene is a member of the heparan sulfate biosynthetic enzyme family. It is a

type II integral membrane protein and possesses heparan sulfate glucosaminyl 3-O-

sulfotransferase activity. The sulfotransferase domain of this enzyme is highly similar to the same domain of heparan sulfate D-glucosaminyl 3-O-sulfotransferase 3B1, and these two enzymes sulfate an identical disaccharide. This gene is widely expressed, with the most

abundant expression in liver and placenta.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

3OST3A1; 30ST3A1; h3-OST-3A; HS3ST3A

## **Product images:**



Western blot analysis of extracts of U-251MG cells, using HS3ST3A1 antibody (TA377226) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.