

## **Product datasheet for TA377255**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **HSPA4L Rabbit Polyclonal Antibody**

**Product data:** 

Product Type:Primary AntibodiesApplications:ELISA, ICC/IF, IF, WBRecommended Dilution:WB,1:500 - 1:2000

IF/ICC,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Formulation:** Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 95kDa

**Gene Name:** heat shock protein family A (Hsp70) member 4 like

**Database Link:** Entrez Gene 22824 Human

<u>095757</u>

**Background:** The protein encoded by this gene is heat shock inducible and may act as a chaperone. The

encoded protein can protect the heat-shocked cell against the harmful effects of aggregated

proteins. This gene is highly expressed in leukemia cells and may be a good target for

therapeutic intervention. Several transcripts encoding different isoforms have been found for


this gene.

**Synonyms:** APG-1; APG1; Osp94





## **Product images:**



Western blot analysis of various lysates using HSPA4L Rabbit pAb (TA377255) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit . Exposure time: 5s.

Immunofluorescence analysis of A431 cells using HSPA4L Rabbit pAb (TA377255) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.