

Product datasheet for TA377284

5HT4 Receptor (HTR4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human HTR4.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 40kDa/42kDa/43kDa/44kDa/45kDa/46kDa/48kDa

Gene Name: 5-hydroxytryptamine receptor 4

Database Link: Entrez Gene 3360 Human

Q13639

Background: This gene is a member of the family of serotonin receptors, which are G protein coupled

receptors that stimulate cAMP production in response to serotonin (5-hydroxytryptamine). The gene product is a glycosylated transmembrane protein that functions in both the

peripheral and central nervous system to modulate the release of various neurotransmitters. Multiple transcript variants encoding proteins with distinct C-terminal sequences have been

described.

Synonyms: 5-HT-4; 5-HT4R




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:





Western blot analysis of extracts of various cell lines, using HTR4 antibody (TA377284) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.

Western blot analysis of extracts of Jurkat cells, using HTR4 antibody (TA377284) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.