

## **Product datasheet for TA377419**

**IL13 Rabbit Polyclonal Antibody** 

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 25-146 of

human IL13 (NP\_002179.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 15kDa

**Gene Name:** interleukin 13

Database Link: Entrez Gene 3596 Human

P35225

**Background:** This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells.

This cytokine is involved in several stages of B-cell maturation and differentiation. It upregulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-

inflammatory cytokines and chemokines. This cytokine is found to be critical to the

pathogenesis of allergen-induced asthma but operates through mechanisms independent of

IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on

chromosome 5q, with this gene particularly close to IL4.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ALRH; BHR1; IL-13; MGC116786; MGC116788; MGC116789; NC30; P600

## **Product images:**



Western blot analysis of extracts of various cell lines, using IL13 antibody (TA377419) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 90s.