

## **Product datasheet for TA377487**

### **ILF2 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of

human ILF2 (NP\_004506.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 43kDa

**Gene Name:** interleukin enhancer binding factor 2

Database Link: Entrez Gene 3608 Human

Q12905



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

The protein encoded by this gene is a transcription factor required for T-cell expression of the interleukin 2 gene. It also binds RNA and is an essential component for encapsidation and protein priming of hepatitis B viral polymerase. The encoded 45 kDa protein (NF45, ILF2) forms a complex with the 90 kDa interleukin enhancer-binding factor 3 (NF90, ILF3), and this complex has been shown to affect the redistribution of nuclear mRNA to the cytoplasm, to repair DNA breaks by nonhomologous end joining, and to negatively regulate the microRNA processing pathway. Knockdown of NF45 or NF90 protein retards cell growth, possibly by inhibition of mRNA stabilization. Alternative splicing results in multiple transcript variants. Related pseudogenes have been found on chromosomes 3 and 14.

Synonyms: MGC8391; NF45; PRO3063

# **Product images:**



Western blot analysis of extracts of various cell lines, using ILF2 antibody (TA377487) at 1:1000 dilution.\_Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.\_Lysates/proteins: 25ug per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Basic Kit .\_Exposure time: 300s.