

## **Product datasheet for TA377570**

## **MUM1 (IRF4) Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Recommended Dilution:** WB,1:500 - 1:2000

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

**Clonality:** Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 150-350 of

human IRF4 (NP\_002451.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 51kDa

**Gene Name:** interferon regulatory factor 4

Database Link: Entrez Gene 3662 Human

Q15306

**Background:** The protein encoded by this gene belongs to the IRF (interferon regulatory factor) family of

transcription factors, characterized by an unique tryptophan pentad repeat DNA-binding domain. The IRFs are important in the regulation of interferons in response to infection by virus, and in the regulation of interferon-inducible genes. This family member is lymphocyte specific and negatively regulates Toll-like-receptor (TLR) signaling that is central to the activation of innate and adaptive immune systems. A chromosomal translocation involving

this gene and the IgH locus, t(6;14)(p25;q32), may be a cause of multiple myeloma.

Alternatively spliced transcript variants have been found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: IRF-4; LSIRF; MUM1; NF-EM5

## **Product images:**



Western blot analysis of extracts of various cell lines, using IRF4 antibody (TA377570) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 60s.