

Product datasheet for TA377611

ITGB1BP1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: **IgG**

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of

human ITGB1BP1 (NP_004754.1).

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. Formulation:

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 16kDa/21kDa

Gene Name: integrin subunit beta 1 binding protein 1

Database Link: Entrez Gene 9270 Human

014713

Background: The cytoplasmic domains of integrins are essential for cell adhesion. The protein encoded by

> this gene binds to the beta1 integrin cytoplasmic domain. The interaction between this protein and beta1 integrin is highly specific. Two isoforms of this protein are derived from alternatively spliced transcripts. The shorter form of this protein does not interact with the beta1 integrin cytoplasmic domain. The longer form is a phosphoprotein and the extent of its phosphorylation is regulated by the cell-matrix interaction, suggesting an important role of this protein during integrin-dependent cell adhesion. Several transcript variants, some

protein-coding and some non-protein coding, have been found for this gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

bodenin; DKFZp686K08158; ICAP-1; ICAP-1A; ICAP-1alpha; ICAP-1B; ICAP1; ICAP1A; ICAP1B; OTTHUMP00000195586

Product images:



Western blot analysis of extracts of 22RV1 cells, using ITGB1BP1 antibody (TA377611) at 1:3000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Enhanced Kit. |Exposure time: 3min.