

## **Product datasheet for TA377707**

## Product datasneet for TAS///0/

## **KCNG4** Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of

human KCNG4 (NP\_758857.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 29kDa/58kDa

**Gene Name:** potassium voltage-gated channel modifier subfamily G member 4

Database Link: Entrez Gene 93107 Human

Q8TDN1

**Background:** Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion

channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily G. This member functions as a modulatory subunit. The gene has strong expression in brain. Multiple alternatively spliced variants have been found in normal and cancerous tissues.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

KCNG3; KV6.3; KV6.4; MGC4558; MGC129609

## **Product images:**



Western blot analysis of extracts of various cell lines, using KCNG4 antibody (TA377707) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 60s.