

## **Product datasheet for TA377728**

## **KCNK10 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 384-543 of

human KCNK10 (NP\_612191.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 59kDa/60kDa

Gene Name: potassium two pore domain channel subfamily K member 10

**Database Link:** P57789

Background: The protein encoded by this gene belongs to the family of potassium channel proteins

> containing two pore-forming P domains. This channel is an open rectifier which primarily passes outward current under physiological K+ concentrations, and is stimulated strongly by arachidonic acid and to a lesser degree by membrane stretching, intracellular acidification, and general anaesthetics. Several alternatively spliced transcript variants encoding different

isoforms have been identified for this gene.

FLJ43399; K2p10.1; TREK-2; TREK2 Synonyms:




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of various cell lines, using KCNK10 antibody (TA377728) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 60s.