

Product datasheet for TA377878S

OriGene Technologies, Inc.9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

MLL5 (KMT2E) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:1000

IHC-P,1:50 - 1:200 IF/ICC,1:50 - 1:200

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 205kDa

Gene Name: lysine methyltransferase 2E

Database Link: Entrez Gene 55904 Human

Q8IZD2

Background: This gene is a member of the myeloid/lymphoid or mixed-lineage leukemia (MLL) family and

encodes a protein with an N-terminal PHD zinc finger and a central SET domain.

Overexpression of the protein inhibits cell cycle progression. Alternate transcriptional splice


variants have been characterized.

Synonyms: HDCMC04P; MLL5






Product images:




Western blot analysis of lysates from C2C12 cells




Immunohistochemistry analysis of paraffinembedded Rat lung using MLL5 Rabbit pAb ([TA377878]) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.





Immunohistochemistry analysis of paraffinembedded Rat spleen using MLL5 Rabbit pAb ([TA377878]) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of paraffinembedded mouse brain using MLL5 Rabbit pAb ([TA377878]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.