

## **Product datasheet for TA377917**

## KRT81 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse

Modifications: Unmodified

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 240-390 of

human KRT81 (NP\_002272.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 54kDa

Gene Name: keratin 81

Database Link: Entrez Gene 3887 Human

Q14533

**Background:** The protein encoded by this gene is a member of the keratin gene family. As a type II hair

keratin, it is a basic protein which heterodimerizes with type I keratins to form hair and nails. The type II hair keratins are clustered in a region of chromosome 12q13 and are grouped into two distinct subfamilies based on structure similarity. One subfamily, consisting of KRTHB1, KRTHB3, and KRTHB6, is highly related. The other less-related subfamily includes KRTHB2, KRTHB4, and KRTHB5. All hair keratins are expressed in the hair follicle; this hair keratin, as well as KRTHB3 and KRTHB6, is found primarily in the hair cortex. Mutations in this gene and KRTHB6 have been observed in patients with a rare dominant hair disease, monilethrix.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ghHb1; ghHkb1; Hb-1; HB1; hHAKB2-1; K2.9; K81; Keratin-81; KRTHB1; MLN137

## **Product images:**



Western blot analysis of extracts of various cell lines, using KRT81 antibody (TA377917) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.