

## **Product datasheet for TA377989**

## **LEFTY2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB.1:500 - 1:2000 Reactivity: Human, Mouse, Rat

**Modifications:** Unmodified

Host: Rabbit Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 57-366 of

human LEFTY2 (NP\_003231.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 37kDa/40kDa

Gene Name: left-right determination factor 2

Database Link: Entrez Gene 7044 Human

O00292

Background: This gene encodes a member of the TGF-beta family of proteins. The encoded protein is

> secreted and plays a role in left-right asymmetry determination of organ systems during development. The protein may also play a role in endometrial bleeding. Mutations in this gene have been associated with left-right axis malformations, particularly in the heart and lungs. Some types of infertility have been associated with dysregulated expression of this gene in the endometrium. Alternative processing of this protein can yield three different products. This gene is closely linked to both a related family member and a related pseudogene. Alternate splicing of this gene results in multiple transcript variants.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

EBAF; LEFTA; LEFTYA; MGC46222; TGF-beta-4; TGFB4

## **Product images:**



Western blot analysis of extracts of various cell lines, using LEFTY2 antibody (TA377989) at 1:1000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 5s.