

# **Product datasheet for TA378052S**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# LIMS1 Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ELISA, ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF/ICC,1:50 - 1:100

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 37kDa

**Gene Name:** LIM zinc finger domain containing 1

**Database Link:** Entrez Gene 3987 Human

P48059






### Background:

The protein encoded by this gene is an adaptor protein which contains five LIM domains, or double zinc fingers. The protein is likely involved in integrin signaling through its LIM domain-mediated interaction with integrin-linked kinase, found in focal adhesion plaques. It is also thought to act as a bridge linking integrin-linked kinase to NCK adaptor protein 2, which is involved in growth factor receptor kinase signaling pathways. Its localization to the periphery of spreading cells also suggests that this protein may play a role in integrin-mediated cell adhesion or spreading. Several transcript variants encoding different isoforms have been found for this gene.


Synonyms:

OTTHUMP00000161608; PINCH; PINCH-1; PINCH1

## **Product images:**



Western blot analysis of lysates from HepG2 cells



Immunofluorescence analysis of MCF7 cells using LIMS1 Rabbit pAb ([TA378052]). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.