

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA378066

Hormone sensitive lipase (LIPE) Rabbit Polyclonal Antibody


Product data:

| Product Type: | Primary Antibodies |
|-------------------------|---|
| | - |
| Applications: | WB |
| Recommended Dilution: | WB,1:500 - 1:2000 |
| Reactivity: | Human, Mouse, Rat |
| Modifications: | Phospho S563 |
| Host: | Rabbit |
| lsotype: | lgG |
| Clonality: | Polyclonal |
| Immunogen: | A phospho specific peptide corresponding to residues surrounding S853 of human HSL (NP_005348.2 ,equivalent to S563 of rat HSL |
| Formulation: | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Shelf life: one year from despatch. |
| Predicted Protein Size: | 84kDa/116kDa |
| Gene Name: | lipase E, hormone sensitive type |
| Database Link: | <u>Entrez Gene 3991 Human</u> <u>Q05469</u> |
| Background: | The protein encoded by this gene has a long and a short form, generated by use of alternative translational start codons. The long form is expressed in steroidogenic tissues such as testis, where it converts cholesteryl esters to free cholesterol for steroid hormone production. The short form is expressed in adipose tissue, among others, where it hydrolyzes stored triglycerides to free fatty acids. |
| Synonyms: | HSL; LHS |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Western blot analysis of extracts of various cell lines, using Phospho-HSL-S563 antibody (TA378066) at 1:1000 dilution.3T3L1 cells were treated by λ -PP mixed solution (1ul) at 30°C for 30 minutes.H9C2 cells were treated with Isoproterenol (10 μ M for 20 min) at 37°C.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per Iane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit .|Exposure time: 180s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US