

Product datasheet for TA378127

LRP6 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Phospho S1490

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic phosphorylated peptide around S1490 of human LRP6.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 180kDa

Gene Name: LDL receptor related protein 6

Database Link: Entrez Gene 4040 Human

<u>075581</u>

Background: This gene encodes a member of the low density lipoprotein (LDL) receptor gene family. LDL

receptors are transmembrane cell surface proteins involved in receptor-mediated

endocytosis of lipoprotein and protein ligands. The protein encoded by this gene functions as

a receptor or, with Frizzled, a co-receptor for Wnt and thereby transmits the canonical Wnt/beta-catenin signaling cascade. Through its interaction with the Wnt/beta-catenin signaling cascade this gene plays a role in the regulation of cell differentiation, proliferation, and migration and the development of many cancer types. This protein undergoes gamma-secretase dependent RIP- (regulated intramembrane proteolysis) processing but the precise

locations of the cleavage sites have not been determined.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ADCAD2; FLJ90062; FLJ90421

Product images:



Western blot analysis of extracts of HeLa cells, using Phospho-LRP6-S1490 antibody (TA378127) at 1:500 dilution.HeLa cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 90s.